

REVIEWARTICLE**5-HT_{2A} Receptor: A Newer Target for Obesity****Jignesh Patel^{1*}, Krupali Mistry², Anirudh Jain³**

- 1. Department of Pharmaceutics and Pharmaceutical Technology, SAL Institute of Pharmacy, Gujarat, India**
- 2. Drug Inspector, Food and drug department, Gandhinagar, Gujarat, India.**
- 3. Department of Pharmaceutics and Pharmaceutical Technology, SAL Institute of Pharmacy, Gujarat, India.**

**Article Received on
20 June 2013.****Revised on 30 June 2013****Accepted on 10 July 2013*****Correspondence for
Author:****Mr. Jignesh Patel**

Asst. Professor,
SAL Institute of Pharmacy
Department of Pharmaceutics
and Pharmaceutical Technology
Opp of Science City, Village:
Bhadaj, Ahmedabad-380060,
Gujarat, India
E mail: jigneshp712@gmail.com

ABSTRACT

Obesity has become major worldwide health problems. Serotonin (5-hydroxytryptamine; 5-HT) is a neurotransmitter that contributes to the regulation of many physiological processes and abnormalities of the serotonergic system have been implicated in the pathogenesis of obesity. 5-HT_{2A} receptor is belongs to G-protein coupled receptor (GPCR), expressed widely throughout the central nervous system (CNS). Hypothalamic 5-HT_{2A} receptors might have a role in the regulation of feeding and energy homeostasis. 5-HT_{2A} receptor gene expression was increased in association with obesity. 5-HT_{2A} receptor antagonism increases expression of adiponectin and decreases plasminogen activator inhibitor 1 (PAI-1) expression via the 5-HT_{2A} receptor signaling cascade. Recently, development of 5-HT_{2A} receptor antagonists as a novel therapeutic strategy for obesity and associated comorbidities has been the focus of much interest. Here, we describe the role of 5-HT_{2A} receptor in pathogenesis of obesity.

Key words: Obesity, serotonin, 5-HT_{2A} receptor, sarpogrelate, adiponectin, PAI-1

INTRODUCTION

Obesity is a medical condition in which excess body fat has accumulated to the extent that it may have an adverse effect on health, leading to reduced life expectancy and/or increased health problems¹. Diet-induced obesity and the subsequent development of features of the metabolic syndrome have become major worldwide health problems. Almost 70% of adults in U.S.A. are overweight but, perhaps more alarmingly, 16% of juveniles are overweight². The number of overweight individuals worldwide has reached 2.1 billion, leading to an explosion of obesity-related health problems associated with a high mortality rate³. Obesity is considered central to the metabolic syndrome and is associated with increases in the risk of an array of diseases, including insulin resistance, type 2 diabetes mellitus, fatty liver disease, atherosclerosis, cardiovascular disease, degenerative disorders, and some cancers⁴. Given that attempts to regulate food intake and content are futile in most of the at risk patients, a clearer understanding of the cellular events underlying the pathophysiology of the obesity is required to allow therapeutic synergism of novel medications along with diet and exercise⁵. Here, we try to emphasize on molecular regulation of 5-HT_{2A} receptor in pathogenesis of obesity.

Serotonin (5-hydroxytryptamine; 5-HT) is a neurotransmitter that contributes to the regulation of many physiological processes such as sleep, appetite, and hormone secretion⁶. Abnormalities of the serotonergic system have been implicated in a number of human diseases including obesity⁷. Acute administration of serotonergic compounds altered the expression of peptidergic appetitive effectors within the hypothalamus, namely pro-opiomelanocortin (POMC) and neuropeptide Y (NPY). POMC and NPY are synthesized within discrete neuronal populations of the arcuate nucleus (ARC) of hypothalamus. Administration of serotonergic compounds causes an increase in anorectic POMC mRNA and a decrease in orexigenic NPY mRNA in arcuate nucleus⁸. Recently, it has been shown that manipulation of these first order hypothalamic POMC/cocaine and amphetamine regulated transcript (CART) and agouti-related protein (AgRP)/NPY neurons is a mechanism through which serotonergic compounds reduce food intake⁹. Specifically, the serotonin system concomitantly regulates the antagonistic functions of POMC/CART and AgRP/NPY neurons through two distinct G-protein coupled receptor subtypes: G_q-coupled 5-HT_{2C} receptor and G_i-coupled 5-HT_{1B} receptor¹⁰. 5-HT_{2C} receptor depolarizes anorectic POMC/CART neurons and release α -melanocyte-stimulating hormone (α -MSH), which in turn activates second-order melanocortin 4 receptor (MC4R) of paraventricular nucleus of the hypothalamus¹¹. Concomitant activation of 5-HT_{1B} receptors expressed on orexigenic AgRP/NPY neurons within the ARC causes membrane hyperpolarization and subsequent

inhibition of neuropeptide release¹². Inhibitory 5-HT_{1B} receptor activation also attenuates inhibitory postsynaptic currents onto POMC/CART neurons and thereby further potentiating anorexigenesis¹³.

Infusion of serotonin into the paraventricular hypothalamus (PVH) of rats reduces food intake¹⁴. Corticotropin-releasing hormone (CRH) neurons located within the PVH have been reported to express MC4R which is responsible for decrease in food intake¹⁵. CRH are directly innervated by serotonergic projections and CRH expression are stimulated by compounds increasing serotonergic efficacy¹⁶. It is possible that serotonin may directly influence the activity of these CRH MC4R-expressing cells and thereby reduces food intake. Thus, an increase in serotonin bioavailability (due to food intake or pharmacological compounds such as sibutramine and fenfluramine) or direct agonism of 5-HT_{2C} receptors and 5-HT_{1B} receptors modulates firing of POMC/CART neurons as well as AgRP/NPY neurons within the arcuate nucleus of the hypothalamus and thereby promotes satiety and the cessation of food intake¹⁷ (Figure I).

Over the years, seven classes of serotonin (5-HT) receptors (5-HT₁ to 5-HT₇) have been identified that are divided into 14 subfamilies. The 5-HT₂ class includes three subtypes of G-protein-coupled receptors, classified as 5-HT_{2A}, 5-HT_{2B} and 5-HT_{2C}¹⁸.




Figure I. Proposed model of a serotonergic pathway modulating food intake¹⁷.

5-HT_{2A} receptors are expressed widely throughout the central nervous system (CNS) especially in cortex (mainly prefrontal, parietal, and somatosensory cortex), olfactory tubercle, midbrain, and cerebellum¹⁹. The high concentrations of 5-HT_{2A} receptors on the apical dendrites of pyramidal cells in layer V of the cortex may modulate cognitive processes, by enhancing glutamate release followed by a complex range of interactions with the 5-HT_{1A}, GABA_A, adenosine A₁, AMPA, mGluR_{2/3}, mGlu₅, and OX₂ receptors²⁰. The 5-HT_{2A} receptors have also been found in the Golgi cells of the granular layer, and in the Purkinje cells of cerebellum²¹. In the periphery, it is highly expressed in platelets and many cell types of the cardiovascular system, in fibroblasts, and in neurons of the peripheral nervous system²².

The 5-HT_{2A} receptor is coded by the *HTR2A* gene. In humans, 5-HT_{2A} gene is located on 13q14-q21 on chromosome 13 and consists of three exons separated by two introns and spans over 20 kb²². More recent data suggest that the *MspI* polymorphism of the 5-HT_{2A} gene may influence food and alcohol intake in obese subjects²³. Some studies have also been indicated a role of the -1438G/A variant of the 5-HT_{2A} gene in the pathogenesis of anorexia nervosa²⁴. The -1438G/A promoter variant is also involved in the pathogenesis of abdominal obesity & related perturbations in insulin, glucose and lipid metabolism as well as in regulation of circulating hormones including salivary cortisol²⁵.

5-HT_{2A} receptors that belong to the super family of G-protein coupled receptors (Gαq-coupled receptors)²⁶. The 5-HT_{2A} receptors activate the phosphoinositide hydrolysis signaling cascade, leading to neuronal depolarization and increases in excitability²⁷. Upon receptor stimulation with agonist, Gα_q and β-γ subunits dissociate to initiate downstream effector pathways. Gα_q stimulates phospholipase C (PLC) activity, results in phospholipase C (PLC)-mediated phosphatidylinositol (PI) lipid hydrolysis, which liberates the second messengers diacylglycerol (DAG) and inositol triphosphates (IP₃), which in turn stimulate protein kinase C (PKC) activity and Ca²⁺ release²⁸. They share a high degree of amino acid sequence homology (68–79% in the transmembrane segments) and similar pharmacological profiles and signal transduction systems with other 5-HT₂ receptor subtypes (5-HT_{2B} and 5-HT_{2C} receptors)²⁹. Activation of 5-HT_{2A} excites GABAergic interneurons in the dorsal raphe nucleus, leading to inhibition of serotonergic cell firing³⁰.

Ligands of 5-HT_{2A} receptor include LSD, psilocin and mescaline act as full or partial agonists at 5-HT_{2A} receptor, and represent the three main classes of 5-HT_{2A} agonists, the ergolines,

tryptamines and phenethylamines, respectively³¹. Ketanserin is a 5-HT_{2A} receptor antagonist with α_1 -adrenoceptor blocking property³².

Sarpogrelate is specific 5-HT_{2A} receptor antagonist and has only insignificant 5-HT₁, 5-HT₃, 5-HT₄, α_1 -adrenoceptor, α_2 -adrenoceptor, α -adrenoceptor, H₁ and H₂-histaminic receptor, and M₃-muscarinic receptor antagonistic activities³³.

5-HT induced platelet activation and platelet aggregation is mediated by 5-HT_{2A} receptor activation³⁴. Acceleration of 5-HT mediated platelet activation at the site of vascular injury and vascular smooth muscle cell proliferation by 5-HT_{2A} receptor activation leads to vascular occlusion. 5-HT_{2A} receptor activation is also involved in the 5-HT-mediated increase in Ca²⁺_i and cause contraction of vascular smooth muscle cells³⁵. Thrombotic and vasoconstrictor effects of 5-HT are mediated by 5-HT_{2A} receptor activation³⁶. Thus, 5-HT_{2A} receptor is of significant clinical interest because of their potential involvement in mediating many cardiovascular diseases³⁷. As 5-HT_{2A} receptor is involved in numerous physiological functions and pathological conditions, it is possible that activating mutations of the 5-HT_{2A} receptor might be responsible for mediating several pathophysiological effects in both the central and peripheral nervous systems³⁸.

5-HT_{2A} Receptor in Obesity

Hypothalamic 5-HT_{2A} receptors might have a role in the regulation of feeding and energy homeostasis. 5-HT_{2A} receptors are likely to down-regulate POMC, CART, CRH, 5-HT_{2C}, and 5-HT_{1B} receptor gene expression in the hypothalamus³⁹.

Hypothalamic 5-HT_{2A} receptor gene expression was increased in association with obesity in A^y mice compared with wild type mice⁴⁰. A^y mice have dominant alleles at the agouti locus (A), and display hyperphagia and obesity⁴¹. It was reported that pharmacologic inactivation of 5-HT_{2A} receptors suppressed hyperphagia and body weight gain, leading to decreased blood glucose levels in obese A^y mice⁴². Hypothalamic 5-HT_{2A} receptors might therefore be involved in the development of obesity and diabetes in A^y mice⁴³. Sarpogrelate, a 5-HT_{2A} receptor antagonist inactivates 5-HT_{2A} receptors and interacts with POMC neurons in the hypothalamus. It stimulates POMC neurons to release enough α -MSH to overcome agouti blockade of MC receptors⁴⁴.

Inhibition of 5-HT might improve insulin sensitivity in diabetes. 5-HT_{2A} receptor mediates hyperglycemic effects of 5-HT through the release of adrenaline from adrenal gland. Adrenaline increases hepatic glucose production and inhibits insulin secretion and the glucose uptake by tissue⁴⁵. Increase in plasma 5-HT level as well as increase in 5-HT release from platelet was

observed in the diabetic patients. These will lead to increase sensitivity to 5-HT in diabetes and hyperglycemia will occur. Therefore, it is thought that inhibition of 5-HT_{2A} might improve insulin sensitivity and thereby led to improvement of insulin resistance⁴⁶.

Adipose tissue participates in the regulation of energy homeostasis, immune responses, and hemostasis as an important endocrine organ that secretes adipokines⁴⁷. In obesity, hypertrophic adipocytes decrease expression and secretion of adiponectin⁴⁸. Adiponectin is an anti-diabetic and anti-atherogenic adipokine⁴⁹. Human adipose tissue contributes to the elevation of plasma plasminogen activator inhibitor 1 (PAI-1) concentrations. PAI-1 plays important roles in the pathogenesis of cardiovascular events, promoting both thrombosis and fibrosis⁵⁰. Among the active 5-HT receptors (1A, 1B, 1D, and 2A), the 5-HT_{2A} receptor was more abundant in hypertrophic adipocytes⁵¹. Expression of 5-HT_{2A} receptor mRNA was increased in hypertrophic 3T3-L1 adipocytes and in mesenteric adipose tissue of diabetic-obese mice, db/db mice, which exhibit decreased expression of adiponectin and increased expression of PAI-1⁵². There is the involvement of the 5-HT_{2A} receptor signaling cascade via mitogen-activated protein kinase (MAPK)-dependent pathways in the regulation of adiponectin and PAI-1 expression⁵³.

Knowledge of the regulatory factors associated with down-regulation of adiponectin gene expression and up-regulation of PAI-1 gene expression is crucial for understanding the pathophysiological basis of obesity and metabolic diseases and could establish new treatment strategies for these conditions⁵⁴. Adiponectin has insulin-sensitizing actions and obesity decreases adiponectin sensitivity, thereby leading to insulin resistance, which in turn aggravates hyperinsulinemia⁵⁵. Expression of PAI-1 was increased in hypertrophic 3T3-L1 adipocytes, which produced a decrease in adiponectin expression. These results are consistent with the adipocyte dysfunction shown in obesity and type2 diabetes⁵⁶. Sarpogrelate increases adiponectin expression⁵⁷. This augmentation was inhibited by suppression of the 5-HT_{2A} receptor gene using siRNA and suppression of this gene also increased adiponectin expression⁵⁸. 5-HT_{2A} stimulation activates Gq protein coupled to the 5-HT_{2A} receptor, decreased adiponectin expression. These findings indicate that the 5-HT_{2A} receptor signaling cascade negatively regulates adiponectin expression⁵⁹. Moreover, expression of the 5-HT_{2A} receptor was up-regulated in the adipose tissue of db/db mice and 3T3-L1 hypertrophic adipocytes, in which adiponectin expression was down-regulated and PAI-1 expression was up-regulated. So, there is possibility that the increase in 5-HT_{2A} receptor expression in hypertrophic adipocytes is at least partially responsible for the

obesity-linked reduction in adiponectin expression. Long-lasting 5-HT_{2A} receptor blockade might increase adiponectin expression down-regulated in obesity⁶⁰. Transcriptional activity of PPAR gamma which increases adiponectin levels has been reported to decrease by MAPK phosphorylation⁶¹. 5-HT_{2A} receptor stimulates MAPK in pulmonary artery fibroblasts which cause proliferative signals. 5-HT_{2A} receptor stimulation may decrease the expression of adiponectin by reduction in the transcriptional activity of PPAR gamma through activation of MAPK in adipocytes⁶². The 5-HT_{2A} receptor signaling cascade could modulate PAI-1 expression through MAPK pathway activation in adipocytes. Arrestin binds to the 5-HT_{2A} receptor. It has been reported that arrestin binding to GPCR enables MAPK activation which is related to increase in PAI-1 gene expression in kidney⁶³.

In summary, 5-HT_{2A} receptor gene expression was increased in association with obesity. 5-HT_{2A} receptor antagonism increases expression of adiponectin and decreases PAI-1 expression via the 5-HT_{2A} receptor signaling cascade. Antagonism of 5-HT_{2A} receptors has the potential to protect against risk factors for and contribute to the treatment of cardiovascular diseases associated with metabolic syndrome as a result of obesity-related, aberrant adipocytokine metabolism. Additional research is required to determine the more precise role of 5-HT_{2A} receptor in obesity and related complications. Such further research investigating the down- and upstream pathways through which serotonin influences appetite may yield additional pharmacological targets for the treatment of obesity.

REFERENCES:

1. Haslam DW, James WP: Obesity. *Lancet*, 2005, 366, 1197–1209.
2. Reaven GM: The insulin resistance syndrome: definition and dietary approaches to treatment. *Annu Rev Nutr*, 2005, 25, 391–406.
3. Sebekova K, Raucinova M, Dzurik R: Serotonin metabolism in patients with decreased renal function. *Nephron*, 1989, 53, 229–232.
4. Huang XF, Han M, Huang X, Zavitsanou K, Deng C: Olanzapine differentially affects 5-HT_{2A} and 5-HT_{2C} receptor mRNA expression in the rat brain. *Behav Brain Res*, 2006, 171, 355–362.
5. Vitale ML, Chiochio SR: Serotonin, a neurotransmitter involved in the regulation of luteinizing hormone release. *Endocr Rev*, 1993, 14, 480–493.
6. Haider S, Haleem DJ: Decreases of brain serotonin following a food restriction schedule of 4 weeks in male and female rats. *Med Sci Monit*, 2000, 6, 1061–1067.

7. Haleem DJ, Haider S: Food restriction decreases serotonin and its synthesis rate in the hypothalamus. *Neuroreport*, 1996, 7, 1153–1156.
8. Boston BA, Blaydon KM, Varnerin J, Cone RD: Independent and additive effects of central POMC and leptin pathways on murine obesity. *Science*, 1997, 278, 1641–1644.
9. Elias CF, Lee CE, Kelly JF, Ahima RS, Kuhar M, Saper CB, Elmquist JK: Characterization of CART neurons in the rat and human hypothalamus. *J Com Neurol*, 2001, 432, 1–19.
10. Seuwen K, Magnaldo I, Pouyssegur J: Serotonin stimulates DNA synthesis in fibroblasts acting through 5-HT_{1B} receptors coupled to a Gi-protein. *Nature*, 1988, 335, 254–256.
11. Sharma SK, Takeda N, Arneja AS, Dhalla NS: Sarpogrelate inhibits genes involved in vascular neointimal hyperplasia and remodeling. *Atheroscler Hypertens diab*, 2003, 98, 175–186.
12. Lucas JL, Yamamoto A, Hen R: Absence of fenfluramine-induced anorexia and reduced c-fos induction in the hypothalamus and central amygdaloid complex of serotonin 1B receptor knock-out mice. *J Neurosci*, 1998, 18, 5537–5544.
13. Sharma SK, Zahradka P, Chapman D, Kumamoto H, Takeda N, Dhalla NS: Inhibition of serotonin-induced vascular smooth muscle cell proliferation by sarpogrelate. *J Pharmacol Exp Ther*, 1999, 290: 1475–1481.
14. Shibano T, Tanaka T, Morishima Y, Yasuoka M, Fujii F: Pharmacological profile of a new 5-hydroxytryptamine 2 receptor antagonist, DV-7028. *Arch Int Pharmacodyn Ther*, 1992, 319, 114–128.
15. Stoving RK, Hangaard J, Hansen-Nord M, Hagen C: A review of endocrine changes in anorexia nervosa. *J Psychiatr Res*, 2000, 33, 139–152.
16. Leibowitz SF: Brain monoamines and peptides: role in the control of eating behavior. *Fed Proc*, 1986, 45, 1396–1403.
17. Garfield AS, Heisler LK: Pharmacological targeting of the serotonergic system for the treatment of obesity. *J Physiol*, 2009, 587, 49–60.
18. Saxena PR: Serotonin receptors: Subtypes, functional responses and therapeutic relevance. *Pharmacol Ther*, 1995, 66, 339–368.
19. Takada Y, Takada A, Urano T: MCI-9042, the new selective antagonist of serotonergic (5-HT_{2A}) receptors. *Cardiovasc Drug Rev*, 1997, 15, 101–121.

20. Shimizu Y, Minatoguchi S, Hashimoto K: The role of serotonin in ischemic cellular damage and the infarct size-reducing effect of sarpogrelate, a 5-hydroxytryptamine-2 receptor blocker, in rabbit hearts. *J Am Coll Cardiol*, 2002, 40, 1347–1355.
21. Halford JC, Blundell JE: Separate systems for serotonin and leptin in appetite control. *Ann Med*, 2000, 32, 222–232.
22. Shouzu A, Nomura S, Hayakawa T: Effect of sarpogrelate hydrochloride on platelet-derived microparticles and various soluble adhesion molecules in diabetes mellitus. *Clin Appl Thromb Hemost*, 2000, 6, 139–143.
23. Sikorski BW, Hodgson WC, King RG: Thromboxane A₂ receptor stimulation similarly potentiates pressor responses to 5-hydroxytryptamine in perfused hindquarters of non-diabetic and alloxan diabetic rats. *Clin Exp Pharmacol Physiol*, 1991, 18, 237–244.
24. Simpson PJ, Schelm JA, Jakubowski JA, Smallwood JK: The role of serotonin (5HT₂) receptor blockade in myocardial reperfusion injury: Effects of LY53857 in a canine model of myocardial infarction. *J Pharmacol Exp Ther*, 1991, 258: 979–985.
25. Smulders YM: Pathophysiology and treatment of haemodynamic instability in acute pulmonary embolism: The pivotal role of pulmonary vasoconstriction. *Cardiovasc Res*, 2000, 48, 23–33.
26. Stassen FR, Fazzi GE, Leenders PJ, Smits JF: Coronary arterial hyperreactivity and mesenteric arterial hyporeactivity after myocardial infarction in the rat. *J Cardiovasc Pharmacol*, 1997, 29, 780–788.
27. Steiner M, Reinhardt KM, Krammer B, Ernst B, Blann AD: Increased levels of soluble adhesion molecules in type 2 (non-insulin dependent) diabetes mellitus are independent of glycemic control. *Thromb Haemost*, 1994, 72, 979–984.
28. Steyn DW, Odendaal HJ: Randomised controlled trial of ketanserin and aspirin in prevention of pre-eclampsia. *Lancet*, 1997, 350, 1267–1271.
29. Borsini F, Bendotti C, Carli M., Poggesi E, Cohen H: The roles of brain noradrenaline and dopamine in the anorectic activity of diethylpropion in rats: a comparison with d-amphetamine. *Res Commun Chem Patho Pharmacol*, 1992, 26: 3–11.
30. Leibowitz SF, Brown LL: Histochemical and pharmacological analysis of catecholaminergic projections to the perifornical hypothalamus in relation of feeding inhibition. *Brain Res*, 1980, 201, 315–345.
31. Yamanda J, Sugimoto Y, Ujikawa M: The serotonin precursor 5-hydroxytryptophan elevates serum leptin levels in mice. *Eur J Pharmacol*, 1999, 383, 49–51.

32. Serruys PW, Klein W, Tijssen JP: Evaluation of ketanserin in the prevention of restenosis after percutaneous transluminal coronary angioplasty. A multicenter randomized double-blind placebo-controlled trial. *Circulation*, 1993, 88, 1588–1601.
33. Shimazu T, Noma M, Saito M: Chronic infusion of norepinephrine into the ventromedial hypothalamus induces obesity in rats. *Brain Res*, 1986, 369, 215–223.
34. Miner LA, Backstrom JR, Sanders-Bush E, Sesack SR: Ultrastructural localization of serotonin 2A receptors in the middle layers of the rat prelimbic prefrontal cortex. *Neuroscience*, 2003, 116, 107–117.
35. Nichols DE, Nichols CD: Serotonin receptors. *Chem Rev*, 2008, 108, 1614–1641.
36. Welsh DJ, Harnett M, MacLean M, Peacock AJ: Proliferation and signaling in fibroblasts: role of 5-hydroxytryptamine 2A receptor and transporter. *Am J Respir Crit Care Med*, 2004, 170, 252–259.
37. Arita Y: Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem Biophys Res Commun*, 1999, 257, 79–83.
38. Collin M, Hakansson OM, Misane I, Ogren SO, Meister BL: Decreased 5-HT transporter mRNA in neurons of the dorsal raphe nucleus and behavioral depression in the obese leptin deficient ob/ob mouse. *Mol Brain Res*, 2000, 81, 51–61.
39. Finn PD, Cunningham MJ, Rickard DG, Clifton DK, Steiner RA: Serotonergic neurons are targets for leptin in the monkey. *J Clin Endocrinol Metab*, 2001, 86, 422–426.
40. Ebihara K, Ogawa Y, Katsuura G, Numata Y, Masuzaki H, Satoh N, Tamaki M: Involvement of agouti related protein an endogenous antagonist of hypothalamic melanocortin receptor in leptin action. *Diabetes*, 1999, 48, 2028–2033.
41. Dryden S, Brown M, King P, Williams G: Decreased plasma leptin levels in lean and obese Zucker rats after treatment with the serotonin reuptake inhibitor fluoxetine. *Horm Metab*, 1999, 31, 363–366.
42. Calapai G, Corica F, Sautebin L, Campo GM, Buemi M, Mauro VN: Leptin increases serotonin turnover by inhibition of brain nitric oxide synthesis. *J Clin Invest*, 1999, 104, 975–982.
43. Nonogaki K, Nozue K, Oka Y: Increased hypothalamic 5-HT_{2A} receptor gene expression and effects of pharmacologic 5-HT_{2A} receptor inactivation in obese A^y mice. *Biochem Biophys Res Commun*, 2006, 351, 1078–1082.

44. Kokubu N: Persistent insulin-sensitizing effects of sarpogrelate hydrochloride, a serotonin 2A receptor antagonist, in patients with peripheral arterial disease. *Circ J*, 2006, 70, 1451–1456.
45. Harris RB, Zhou J, Redmann SM, Smagin GN, Smith SR, Rodgers E, Zachwieja JJ: A leptin dose-response study in obese (ob/ob) and lean mice. *Endocrinology*, 1998, 139, 8–19.
46. Currie PJ, Coscina DV: Metergoline potentiates natural feeding and antagonizes the anorectic action of medial hypothalamic 5-hydroxytryptamine. *Pharmacol Biochem Behav*, 1996, 53, 1023–1028.
47. Kershaw EE, Flier JS: Adipose tissue as an endocrine organ. *J Clin Endocrinol Metab*, 2004, 89, 2548–2556.
48. Kohler HP, Grant PJ: Plasminogen-activator inhibitor type 1 and coronary artery disease. *New Engl J Med*, 2000, 342, 1792–1801.
49. Yamauchi T: Adiponectin stimulates glucose utilization and fatty-acid oxidation by activating AMP-activated protein kinase. *Nat Med*, 2002, 8, 1288–1295.
50. Kawano H: Serotonin induces the expression of tissue factor and plasminogen activator inhibitor-1 in cultured rat aortic endothelial cells. *Blood*, 2001, 97, 1697–1702.
51. Hoyer D, Clarke DE, Fozard JR: International Union of Pharmacology classification of receptors for 5-hydroxytryptamine (Serotonin). *Pharmacol Rev*, 1994, 46, 157–203.
52. Knight G, Boulton AJ, Drury J, Ward JD: Long term glycaemic control by alternative regimens in a feasibility study of continuous subcutaneous insulin infusion. *Diabetes Res*, 1986, 3, 355–358.
53. Goicolea OI, Hernandez CI, Garcia JA: Comparative study between the subcutaneous continuous insulin infusion pump and optimized conventional treatment: effects at 6 months. *Rev Clin Esp*, 1986, 179, 3–7.
54. Olsen T, Richelsen B, Ehlers N, Beck-Nielsen H: Diabetic retinopathy after 3 years treatment with continuous subcutaneous insulin infusion (CSII). *Acta Ophthalmol*, 1987, 65, 185–189.
55. Fang X, Sweeney G: Mechanisms regulating energy metabolism by adiponectin in obesity and diabetes. *Biochem Soc Transac*, 2006, 34, 798-799.
56. Dahl JS, Toriesen P, Hanssen K, Sandvik L, Aagenes O: Increase in insulin antibodies during continuous subcutaneous insulin infusion and multiple-injection therapy in contrast to conventional treatment. *Diabetes*, 1987, 36, 1–5.

Patel et al. 5-HT_{2A} Receptor: A Newer Target for Obesity

57. Edelman E, Walter H, Biermann E, Schleicher E, Bachmann W, Mehnert H: Sustained normoglycemia and remission phase in newly diagnosed type I diabetic subjects: comparison between continuous subcutaneous insulin infusion and conventional therapy during a one year follow-up. *Horm Metab Res*, 1987, 19, 419–421.
58. Kokubu N: Persistent insulin-sensitizing effects of sarpogrelate hydrochloride, a serotonin 2A receptor antagonist, in patients with peripheral arterial disease. *Circ J*, 2006, 70, 1451–1456.
59. Cerrito F, Lazzaro MP, Gaudio E, Arminio P, Aloisi G: 5HT₂-receptors and serotonin release: their role in human platelet aggregation. *Life Sci*, 1993, 53, 209–215.
60. Nomura S, Shouzu A, Omoto S, Nishikawa M, Iwasaka T: 5-HT_{2A} receptor antagonist increases circulating adiponectin in patients with type 2 diabetes. *Blood Coagul Fibrinolysis*, 2005, 16, 423–428.
61. Kadowaki T: The role of PPAR gamma in high-fat diet-induced obesity and insulin resistance. *J Diabetes Complications*, 2002, 16, 41–45.
62. Kadowaki T, Yamauchi T: Adiponectin and adiponectin receptors. *Endocr Rev*, 2005, 26, 439–451.
63. Nerurkar SS: P38 MAPK inhibitors suppress biomarkers of hypertension end-organ damage, osteopontin and plasminogen activator inhibitor-1. *Biomark*, 2007, 12, 87–112.